## DAWSON COLLEGE MATHEMATICS DEPARTMENT

## Final Examination

Mathematics 201-NYC-05 Linear Algebra (Commerce) Instructors: Rodney Acteson, Melanie Beck, Olga Zlotchevskaia Date: Friday, December 14, 2012 Time: 2:00 - 5:00

1. (6 marks) Solve the following system using Gaussian elimination or Gauss-Jordan elimination.

2. (6 marks) Consider the system

For which value(s) of k is the system

- (a) consistent?
- (b) inconsistent?
- 3. (3 marks) Determine whether the following statement is true or false. If the answer is True, justify your answer, if the answer is False, provide an example which shows that the statement is False.

 $(A + B)(A = B) = A^2 = B^2$ ; for all square matrices A and B of the same size.

- 4. (5 marks) Simplify  $(AB)^{-1}(AC)(A^{T}C)^{-1}$ .
- 5. (6 marks) Find A such that

$$(A^{T} \ 8 \ I)^{1} = \begin{bmatrix} h & 2 & 3 \\ 4 & 5 \end{bmatrix}^{1}$$

6. (4 marks) Find elementary matrices  $E_1$  and  $E_2$  which satisfy the following equation.

$$E_2 E_1 \begin{bmatrix} h & 5 & 6 \\ 0 & 1 \end{bmatrix} = I$$
7. (8 marks) Given  $A = \begin{bmatrix} 4 & 4 & 4 \\ 2 & 0 & 1 \\ 1 & 2 & 3 \end{bmatrix}$ ,  $X = \begin{bmatrix} y \\ z \end{bmatrix}$ , and  $B = \begin{bmatrix} 4 \\ 0 \\ 2 \end{bmatrix}$ , nd

- (a) det(*A*)
- (b) adj(*A*)
- (c)  $A^{-1}$  using adj(A)
- (d) The solution to AX = B using  $A^{-1}$
- 8. (5 marks) Using only row operations, nd the determinant of the following matrix A.

## 9. (9 marks) Let A and B be 3 3 matrices with det(AB) = 24 and det(A) = 6. Find

- (a) det( 3A<sup>1</sup>)
- (b) det(BAB)
- (c)  $det(A^T det(A))$

10. (6 marks) Consider the vectors u = (2; 2; 3) and v = (2; 2; 1).

- (a) Find a unit vector perpendicular to both u and v.
- (b) Find the area of the parallelogram de ned by the vectors u and v.
- 11. (4 marks) Consider the vectors u = (2k; 1; k) and v = (4; 2; 2).
  - (a) For which value(s) of k, are the vectors u and v perpendicular?
  - (b) For which value(s) of k, are the vectors u and v parallel?
- 12. **(4 marks)** Find the equation of the line passing through P(3; 1; 2) and perpendicular to the plane  $4x \quad 5y = 2z + 3$ .
- 13. (6 marks) Given the two planes

2(x + 1) + (y + 1) + 3(z + 4) = 0 and x + y + 4z = 2;

- (a) Show that the planes are not parallel.
- (b) Find the line of intersection of the planes.
- 14. (5 marks) Find the point on the line

$$x = 1 + 3t; y = 2 - 2t; z = 1 - t; (-1 < t < 1);$$

that is closest to P(-1;5;9).

15. (6 marks) Find an equation for the plane that contains the point P(2;1;3) and the line

x = 1 + 2t; y = 2 + t; z = 2 - t; (-1 < t < 1):

16. (5 marks) Prove the following identity

(U + kV) V = U  $V_i$ 

where k is a scalar and u and v are vectors in 3-space.

- 17. **(4 marks)** A manufacture is producing bikes and scooters. It takes 6 hours to make a bike and 4 hours to make a scooter. Each bike requires 7 kg of steel and each scooter requires 5 kg of steel. The manufacture has 40 hours available for making bikes and scooters and has 280 kg of steel on hand. The company makes a pro t of \$25 on each bike and \$15 on each scooter. How many bikes and how many scooters should it make in order to maximize he pro t? Set up the linear programming problem as follows **(you are not asked to solve the problem!)**:
  - Set up the filtear programming problem as follows (you are not asked to solve the pre
  - (a) De ne all the variables (using the phrase \the number of").
  - (b) State the objective and identify the objective function (in terms of the variables).
  - (c) State all the constraints (in terms of the variables).
- 18. (8 marks) Solve the following problem.

Maximize subject to
 
$$p = 4x + 2y$$
 $2z$ 
 $x y z$ 
 $20$ 
 $2x + y + 2z$ 
 $70$ 
 $x + z$ 
 $10$ 
 $x' y' z$ 
 $0$ 

Answers:  
1. 
$$x = 8$$
  $\frac{19}{2}t$ ;  $y = 4$   $\frac{5}{2}t$ ;  $z = t$   
2. (a) if  $k = 1$  (b) if  $k \neq 1$   
3. False: example  $A = {\begin{array}{c} h & 1 & 0 \\ 1 & 0 \end{array}} {\begin{array}{c} B & = \\ 0 & 0 \end{array}} {\begin{array}{c} B & - \\ 0 & 0 \end{array}} {\begin{array}{c} A & - \\ 0 & - \\ 0 \end{array}} {\begin{array}{c} A & - \\ 0 & 0 \end{array}} {\begin{array}{c} A & - \\ 0 \end{array}} {\begin{array}$